

เอกสารอ้างอิง

- กฤษณา รุ่งเรืองศักดิ์. (2521). ปฏิบัติการและหลักเบื้องต้นในวิชาชีวเคมี. อัมรินทร์การพิมพ์, กรุงเทพ.
- ดาวลักษณ์ พิมภู. (2538). ชีวเคมีเล่ม 1 โนโลจีชีวภาพ. กรุงเทพ:สำนักพิมพ์ประกายพรีก.
- เดือนพิพพ์ ปิยรัตน์. (2538). การศึกษาสภาวะที่เหมาะสมต่อการผลิตกรดไขมัน โอมegas-3 จากสาหร่ายนำเข้าขนาดเล็ก; คณะวิทยาศาสตร์, มหาวิทยาลัยศรีนครินทรวิโรฒ ประสานมิตร, กรุงเทพ, ประเทศไทย.
- วินัย คงหล้าน, นปป. สมคุลทางเมทานอลิซึมของลิพิดกับการเกิดโรคและการป้องกัน. เมทานอลิซึมของลิพิดกับการป้องกัน. 15 หน้า
- สุนันทา กิจญาเวชน์. (2535). ชีวเคมี 2. กรุงเทพ:สำนักพิมพ์มหาวิทยาลัยรามคำแหง.
- สุพิช ทองรอด. (2535). ความสำคัญของกรดไขมันในอาหารสัตว์น้ำ. วารสารกรมประมง 4(2): 943-950.
- อนุเทพ ภาสุระ. (2541). เอกสารประกอบการสอนวิชา ไมโครโลยี (Mycology). ภาควิชาจุลชีววิทยา คณะวิทยาศาสตร์ มหาวิทยาลัยบูรพา. ชลบุรี:
- อัตนิตร์ อิทธิอาภา. (2541). การหาปริมาณไขมันและเปอร์เซนต์ของกรดไขมันไม่อิ่มตัวกลุ่มโอมegas-3 ในปลาทะเลโดยเทคนิคแก๊สลิคิวิค โตรมาโตกราฟี, คณะวิทยาศาสตร์, มหาวิทยาลัยบูรพา, ชลบุรี.
- Alexopoulos, G.J., Mims, C.W., Blackwell, M. (1996). Introductory Mycology. New York: John Wiley and Son, Inc.
- Atlas, R.M and Bartha, R., 1981 Microbial Ecology : Fundamentals and Applications. Addison-Wesley Publishing company.london. 550 p.
- Bahnweg, G and Sparrow, F.K. (1972). *Aplanochytrium kerguelensis* gen. Spec. nov., a new phycomycete from subantarctic marine waters. *Archiv fur Mikrobiologie*. 81: 45-49.
- Bajpai, P.K., Bajpai, P.K., and Ward, O. (1991a). Optimization of production of docosahexaenoic acid (DHA) by *Thraustochytrium aureum* ATCC 34304. Journal American Oil Chemistry Society 68 (7): 509-513.
- Bajpai, P.K., Bajpai, P.K., and Ward, O. (1991b). Eicosapentaenoic acid (EPA) formation: Comparative studies with *Mortierella* strain and production by *Mortierella elongata*. Mycology Research 95(11):1294-1298.
- Barclay, W. R. and Zeller, S. (1996). Nutritional enhancement of n-3 and n-6 fatty acids in rotifers and *Artemia nauplii* by feeding spray-dried *Schizochytrium* sp. Journal of The World Aquaculture Society 27: 314-322.

- Bell, M.V., Henderson, R.J. and Sargent, J.R. (1986). The role of polyunsaturated fatty acids in fish. Comparative Biochemistry and Physiology, 83: 711-719.
- Bowles, R.D., Hunt, A.E., Duchars, M.G., and Eaton, R.A. (1999). Long-chain n-3 polyunsaturated fatty acid production by member of the marine protistan group the *Thraustochytrid*: Screening of isolate and optimisation of docosahexaenoic acid production. Journal Biotechnology 70(15):193-195.
- Campbell, I. and Duffus, J. H. (eds). 1991 Yeast a practical approach. Internation Russ limited, Oxford. 289 pp.
- Chirala, S. [Http://www.bcm.tmc.edu/biochem/fac/chirala.html](http://www.bcm.tmc.edu/biochem/fac/chirala.html).
- Christie, W.W. (1989). Gas chromatography and lipids: a practical guide. The Oily Press, Ayr, Scotland.
- Dickson, L.G., Galloway, R.A., and Patterson, G.W. (1969). Environmentally-induced changes in the fatty acid. Plant Physiology 44(23):1413-1418.
- Gaertner, A. (1977). Revision of the Thraustochytriaceae (lower marine fungi). I. *Ulkenia* nov. gen., with description of three new species. *Veroffentlichungen des Instituts fur Meeresforschung in Bremerhaven*. 16: 139-157.
- Goldstein, S. and Belsky, M. (1964). Axenic culture studies of a new marine phycomycete possessing an unusual type of asexual reproduction. American Journal of Botany. 51: 72-78.
- Grima, E.M., Sancha, J.A., Camacho, F.G., Medina, A.R., Gimenez, A.G., and Alonso, D.L. (1994). The production of polyunsaturated fatty acid by microalgae: From strain selection to product purification. Process Biochemistry 30(8):711-715.
- Gunstone, F.D. (1996). Fatty acid and lipid chemistry. Blackie Academic & Professional. London. 252 p.
- Honda, D. (2001). Phylogeny and taxonomy of labyrinthulids. Aquabiology, 132, 23 (1): 7-17.
- Hyde, K.D., Pointing, S.B. (2000). The Marine Mycology : A Practical Approach. Hongkong:Fungal Diversity Press.
- Iida, I., et al (1996). Improvement of docosahexaenoic acid production in a culture of *Thraustochytrium aureum* by medium optimization. Journal of Fermentation and Bioengineering 21(1):76-80.
- Jareonkitmongkol, S., Shimizu, S., and Yamada, H. (1992). Fatty acid desaturation- defective mutants of an arachidonic acid producing fungus , *Mortierella alpina*. 1S- 4. Journal of General Microbiology, 14(38):997-1001.

- Jaritkuan, S.; E.B.G. Jones and Bremer, G. (1998). Thraustochytrids as a food source in aquaculture. Paper presented at The Asia-Pacific Mycological Conference on Biodiversity and Biotechnology, 6-9 July, 1998. Hua Hin, Thailand.
- Jaritkuan, S. and E.B.G. Jones. (1999). Aquaculture feed from Thraustochytrids. Paper presented at The 7th International Marine and Freshwater Mycology Symposium. 4-9 July 1999. City University of Hong Kong, Hong Kong.
- Jaritkuan, S. (2002). Thraustochytrids: a new alternative source of fatty acids for aquaculture. In: Fungi in Marine Environments (ed. K.D. Hyde). Fungal Diversity Research Series 7:345-357.
- Kendrick, A. and Ratledge, C. (1992). Lipids of selected molds grown for production of n-3 and n-6 polyunsaturated fatty acids. *Lipids*. 27: 15-20.
- Kobayashi, Y. and Ookubo, M. (1953). Studies on the marine phycomycetes. Bulletin of the National Science Museum, Tokyo. 3: 53-65.
- Leano, E.M. (2001). Straminipilous organisms from fallen mangrove leaves from Panay Island, Philippines. *Fungal Diversity* 6: 75-81.
- Li, Z. Y. and Ward, O. P. (1994). Production of docosahexaenoic acid by *Thraustochytrium roseum*. *Journal of Industrial Microbiology*, 13: 238-241.
- Luthisungneon, P. (1998). Study on morphology, physiology and fatty acid profiles of *Aschersonia* for identification of closely related fungal TAXA (MSc. in Biotechnology Program), King Mongkut's University of Technology, Bangkok, Thailand.
- Mackenzie, D.A., Carter, A.T., Wongwathanarat, P., Eagles, J., Salt, J. and Archer, D.B. 2002. A third fatty acid Δ9-desaturases from *Mortierella alpina* with a different substrate specificity to ole 1 p and ole 2 p *Microbiology* 148:1725-1735
- Mannella, C.A., Frank, J. and Delihas, N. (1987). Interrelatedness of 5S RNA sequences investigated by correspondence analysis. *Journal of Molecular Evolution*, 24: 228-235.
- Nakahara, T., Yokochi, T., Higashihara, T., Tanaka, S., Yaguchi, T., and Honda, D. (1996). Production of docosahexaenoic acid and docosapentaenoic acid by *Schizochytrium* sp. Isolate from Yap Island. *Journal American Oil Chemistry Society* 73(11):1421-1425.
- Otero, A., Garcia, D., Morales, E.D., Aran, J., and Jaime, H. (1997). Manipulation of the biochemical composition of the eicosapentaenoic acid rich microalgae *Isochysis galbana* in semicontinuous culture. *Biotechnology and Applied Biochemistry* 26 (12):171-175.

- Raghuram, S. (1987). Occurrence of the thraustochytrid, *Corallochytrium limacisporum* gen et sp. Nov. in the coral reef lagoons of Lakshadweep Islands in the Arabian Sea. *Botanica Marina*. 30: 83-89.
- Rose, A.H. and Harrison, J.S. (eds.) 1971. The yeast : Physiology and Biochemistry of yeast.: Vol 2. Academic Russ. London. 571 p.
- Sargent, J., Bell, G., McEvoy, L., Tocher, D. and Estevez, A. (1999). Recent developments in the essential fatty acid nutrition of fish. *Aquaculture*, 177: 191-199.
- Shimizu, S., Kawashima, H., Shinmen, Y., Akitomo, K. and Yamada, H. (1988). Production of eicosapentaenoic acid by *Mortierella* fungi. *Journal of the American Oil Chemists Society*, 65: 1455-1459.
- Silva, L.T., Sousa, D.E., Pereira, P.T., Eerrao, A.M., and Roscior, J.C. (1998). Cell fatty acid profiles for the differentiation of *Penicillium* species. *FEMS Microbiology Letter* 164(2):303-309.
- Singh, A. and Ward, O.P. (1996). Production of high yields of docosahexaenoic acid by *Thraustochytrium roseum* ATCC 28210. *Journal of Industrial Microbiology*, 16: 370-373.
- Sparrow, F.K. (1936). Biological observations on the marine fungi of Woods Hole waters. *Biological Bulletin*, Marine Biological Laboratory, Woods Hole, Massachusetts. 70:236-263.
- Stahl, P.D., and Klug, M.J. (1996). Characterization and differentiation of filamentous fungi based on fatty acid composition . *Applied and Environmental Microbiology* 62 (11):4136-4141.
- Uauy-Dagach, R. (1996). Omega-3 PUFA in perinatal nutrition: docosahexaenoic acid (DHA) needs during pregnancy and infancy. The paper presented at the Conference on The Essence of Polyunsaturated Fatty Acids (PUFA) in Food Industry. February 14, 1996, Bangkok, Thailand. p: 24-40.
- Voet, D and Voet, J.G. (1995). *Biochemistry*. 2nd ed. Wiley, New York, Chichester, Brisbane, Toronto and Singapore. 1361 pp.
- Wallage,R.A., King,J.L. and Sander, G.P. 1986. *Biology the Science of life*. 2nd Edition. Scott, Foresman and company. London. 1217 p.
- Yaguchi, T., Tanaka, S., Yokochi, T., Nakahara, T., and Higashihara, T. (1997). Production of high yield of docosahexaenoic acid by *Shizochytrium* sp. strain SR21. *Journal American Oil Chemistry Society* 74(11):1431-1436.

190671

- Yazama, K.; Watanabe, K; Ishikawa, C.; Kondo, K, and Kimura, S. (1992). Production of eicosapentaenoic acid from marine bacteria. In: Industrial applications of single cell oil. P.29-51. (Eds. D.J. Kyle, C. Ratledge). American Oil Chemists Soc., Illinois, USA.
- Yongmanitchai, W. and Ward, O.P. (1989). Omega-3 fatty acids: Alternative sources of production. Process Biochemistry, 117-125.
- Yongmanitchai, W., and Ward, O.P. (1991). Growth of and omega-3 fatty acid production by *Phaeodactylum tricornutum* under different culture condition. Applied and Environmental Microbiology 57 (2):419-421.
- Zelles, L. (1917). Phospholipid fatty acid profiles in selected member of soil microbial communities. Chemosphere 35(1):275-281.