

บรรณานุกรม

- สมทรง เลขคุณ. (2542). ชีวเคมีของวิตามิน (พิมพ์ครั้งที่ 1). กรุงเทพฯ: ศุภวนิชการพิมพ์.
- สุวรรณี ธีรภาพธรรมกุล. (2541). การวิเคราะห์ปริมาณแคนเฟเพอินในการแพะปูรุ่งสำเร็จพร้อมคั่มโดยวิธี HPLC. วารสารกรมวิทยาศาสตร์การแพทย์, 41(3), 317-325.
- Albala-Hurtado, S., Veciana-Nogues, M. T., Izquierdo-Pulido, M., & Marine-Font, A. (1997). Determination of water-soluble vitamins in infant milk by high performance liquid chromatography. *Journal of Chromatography A*, 788, 247-253.
- Aucamp, J. P., Hara, Y., & Apostolidis, Z. (2000). Simultaneous analysis of tea catechins, caffeine, gallic acid, theanine and ascorbic acid by micellar electrokinetic chromatography. *Journal of Chromatography A*, 876, 235-242.
- Boonkerd, S., Detaevernier, M. R., & Michotte, Y. (1994). Use of capillary electrophoresis for the determination of vitamins of the B group in pharmaceutical preparation. *Journal of Chromatography A*, 670, 209-214.
- Cataldi, T. R. I., Nardiello, D., Scrano, L., & Scopa, A. (2002). Assay of riboflavin in sample wines by capillary zone electrophoresis and laser-induced fluorescence detection. *Journal of Agricultural and Food Chemistry*, 50, 6643-6647.
- Chen, G., Ding, X., Cao, Z., & Ye, J. (2000). Determination of melatonin and pyridoxine in pharmaceutical preparation for health-caring purposes by capillary electrophoresis with electrochemical detection. *Analytica Chimica Acta*, 408, 249-256.
- Cho, C. M., Ko, J. H., & Cheong, W. J., (2000). Simultaneous determination of water-soluble vitamins excreted in human urine after eating an overdose of vitamin pills by a HPLC method coupled with a solid phase extraction. *Talanta*, 51, 799-806.
- Conte, E. D., Barry, E. F., & Rubinstein, H. (1996). Determination of caffeine in beverages by capillary zone electrophoresis. *Journal of Chemical Education*, 73 (12), 1169-1170.

- Delgado-Zamarreno, M. M., Gonzalez-Maza, I., Sanchez-Perez, A., & Carabias-Martinez, R. (2002). Separation and simultaneous determination of water-soluble and fat-soluble vitamins by electrokinetic capillary chromatography. *Journal of Chromatography A*, 953, 257-262.
- Fotsing, L., Fillet, M., Chiap, P., Hubert, P., & Crommen, J. (1997). Determination of six water vitamins in a pharmaceutical formulations by capillary electrophoresis. *Journal of Pharmaceutical and Biomedical Analysis*, 15, 1113-1123.
- Fotsing, L., Fillet, M., Chiap, P., Hubert, P., & Crommen, J. (1999). Elimination of adsorption effects in the analysis of water-soluble vitamins in pharmaceutical formulations by capillary electrophoresis. *Journal of Chromatography A*, 853, 391-401.
- Fujiwara, S., Iwase, S., & Honda, S. (1988). Analysis of water-soluble vitamins by micellar electrokinetic capillary chromatography. *Journal of Chromatography*, 447, 133-140.
- Gomis, D. B., Gonzalez, L. L., & Alvarez, D. G. (1999). Micellar electrokinetic capillary chromatography analysis of water-soluble vitamins. *Analytica Chimica Acta*, 396, 55-60.
- Grossman, P. D., & Colburn, J. C. (1992). *Capillary electrophoresis. Theory and practice*. San Diego: Eds., Academic Press.
- Huber, L. (1998). *Validation and qualification in analytical laboratories*. Buffalo Grove, IL: Interpharm Press.
- Jegle, U. (1993). Separation of water-soluble vitamins via high performance capillary electrophoresis. *Journal of Chromatography A*, 652, 495-501.
- Lee, B. L., & Ong, C. N. (2000). Comparative analysis of tea catechins and theaflavins by high-performance liquid chromatography and capillary electrophoresis. *Journal of Chromatography A*, 881, 439-447.

- Markopoulou, C. K., Kagkadis, K. A., & Koundourellis, J. E. (2002). An Optimized method for the simultaneous determination of vitamins B₁, B₆, B₁₂, in multivitamin tablets by high performance liquid chromatography. *Journal of Pharmaceutical and Biomedical Analysis*, 30, 1403-1410.
- Moreno, P., & Salvado, V. (2000). Determination of eight water- and fat-soluble in multi-vitamin pharmaceutical formulations by high-performance liquid chromatography. *Journal of Chromatography A*, 870, 207-215.
- Nishi, H., Tsumagari, N., Kakimoto, T., & Terabe, S. (1989). Separation of water-soluble vitamins by micellar electrokinetic chromatography. *Journal of Chromatography*, 465, 331-343.
- Okamoto, H., Nakajima, T., & Ito, Y. (2002). Simultaneous determination of ingredients in a vitamin-enriched drink by micellar electrokinetic chromatography. *Journal of Pharmaceutical and Biomedical Analysis*, 30, 815-822.
- Quirino, J. P., & Terabe, S. (1999). Electrokinetic chromatography. *Journal of Chromatography A*, 856, 465-482.
- Sanchez, J. M., & Salvado, V. (2002). Comparison of micellar and microemulsion electrokinetic chromatography for the analysis of water- and fat-soluble vitamins. *Journal of Chromatography A*, 950, 241-247.
- Su, S-C., Chou, S-S., Hwang, D-F., Chang, P-C., & Liu, C-H. (2001). Capillary zone electrophoresis and micellar electrokinetic chromatography for determining water-soluble vitamins in commercial capsules and tablets. *Journal of Food Science*, 66(1), 10-14.
- Versari, A., Mattioli, A., Parpinello, G. P., & Galassi, S. (2004). Rapid analysis of ascorbic and isoascorbic acids in fruit juice by capillary electrophoresis. *Food Control*, 15, 355-358.
- Walker, J. C., Zaugg, S. E., & Walker, E. B. (1997). Analysis of beverages by capillary electrophoresis. *Journal of Chromatography A*, 781, 481-485.

Watanabe, T., Nishiyama, R., Yamamoto, A., Nagai, S., & Terabe, S. (1998).

Simultaneous analysis of individual catechins, caffeine, and ascorbic acid in commercial canned green and black teas by micellar electrokinetic chromatography. *Analytical Sciences*, 14, 435-438.

Weinberger, R. (1993). *Practical capillary electrophoresis*. New York: Harcourt Brace Jovanovich.

Zhao, Y., & Lunte, C. E. (1997). Determination of caffeine and its metabolites by micellar electrokinetic capillary chromatography. *Journal of Chromatography B*, 688, 265-274.